

: رياضيات 6 (نظرية الزمرة)

: كيمياء (تربية عام)

6 :

: د. هدير الجندی

## Examples of subgroups

Def.

For any element  $a$  from a group,  
we let  $\langle a \rangle$  denote the set

$$\{a^n \mid n \in \mathbb{Z}\}$$

Theorem

Let  $G$  be a group, and let  
 $a$  be any element of  $G$ . Then

$\langle a \rangle$  is a subgroup of  $G$ .

Proof

Since  $a \in \langle a \rangle$ ,  $\langle a \rangle$  is not empty.

Let  $a^n, a^m \in \langle a \rangle$ . Then  $a^n (a^m)^{-1} = a^{n-m} \in \langle a \rangle$ .

Hence  $\langle a \rangle$  is a subgroup of  $G$ .

## Remarks

- (1) The subgroup  $\langle a \rangle$  is called the cyclic subgroup of  $G$  generated by  $a$ .
- (2) In the case that  $G = \langle a \rangle$ , we say that  $G$  is cyclic and  $a$  is a generator of  $G$ .
- (3) Although the list  $\dots, a^{-2}, a^{-1}, a^0, a^1, a^2, \dots$  has infinitely many entries, the set  $\{a^n \mid n \in \mathbb{Z}\}$  might have only finitely many elements.

(4) Note that, since  $a^i a^j = a^{i+j} = a^{j+i} = a^j a^i$ ,

every cyclic group is Abelian.

Example in  $\mathbb{Z}_{10}$ ,  $\langle 2 \rangle = \{2, 4, 6, 8, 10\}$ .